

TOROCITY™



ACTIVE INGREDIENT:	(% by weight)
Mesotrione: 2-[4-(methylsulfonyl)-2-nitrobenzoyl]-1,3-cyclohexanedione	40.0%
OTHER INGREDIENTS:	<u>60.0%</u>
TOTAL:	<u>100.0%</u>

Contains 4 lbs. Mesotrione per gallon.
EPA Reg. No.: 91234-75

SPECIMEN

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See below for additional precautionary statements.

FIRST AID	
If on skin:	<ul style="list-style-type: none"> ▪ Take off contaminated clothing. ▪ Rinse skin immediately with plenty of water for 15 - 20 minutes. ▪ Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	

For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC Day or Night Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with skin, eyes, or clothing.

Personal Protection Equipment (PPE)

Applicators and Other Handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove and wash contaminated clothing before reuse.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

Surface Water Advisory

This product may contaminate water through drift or spray in wind. This product has a high potential for runoff for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

Physical and Chemical Hazards

Do not use or store near heat or open flame. Do not use with or store near any oxidizing or reducing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- coveralls
- shoes plus socks
- chemical-resistant gloves

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard, 40 CFR Part 170. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, and greenhouses.

Do not enter treated areas without protection clothing until sprays have dried.

Torocity is applied pre- and post-emergence to provide selective contact and residual control of turfgrass weeds. If applied pre-emergence, it is absorbed when emerging from the soil. Pre-emergence activity and control is reduced under dry conditions. Activate **Torocity** with 0.15 inches of irrigation if rain hasn't fallen within 10 days of application. Post-emergent control is obtained by absorption into the soil and contact with foliage. Growth ceases post-application, weeds turn white from chlorophyll loss, and will die within three weeks. Make a repeat application after 2 - 3 weeks to improve post-emergence weed control. Add a non-ionic surfactant when making post-emergence applications.

Turfgrass color can temporarily become white during treatment. Whitening typically occurs 5 - 7 days post-application and lasts for several weeks. A second application to the same site will cause less whitening of plant tissue.

Torocity controls weeds prior to and during seeding of certain turfgrasses during turf renovation (see **New Seedings**). If making pre-emergence application to established turf, tank mix **Torocity** with other pre-emergence herbicides such as Barricade® 65WG (EPA Reg. No. 100-834, prodiamine) for longer residual and broad spectrum control.

MANDATORY SPRAY DRIFT MANAGEMENT

Ground Applications

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For all applications, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

Spray Drift Advisories

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray

drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

BOOM HEIGHT - Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

WINDBLOWN SOIL PARTICLES

Torocity has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other factors which can affect the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying Torocity if prevailing local conditions may be expected to result in off-site movement.

APPROVED USE SITES

Torocity can be applied to commercial and residential turfgrasses. Non-crop area use sites include golf courses, sod farms, athletic fields, parks, residential and commercial properties, cemeteries, airports, and lawns.

Turfgrass Use Precautions:

Apply Torocity at reduced rates of 4 fl. oz./A (0.125 lb. AI/A) or less if tank mixing with atrazine, bentazon, or simazine. Before tank mixing Torocity with other herbicides, conduct a compatibility, safety, and efficacy test before treating larger areas. See tank mix partner labels for directions and precautions. The most restrictive directions apply. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Thoroughly clean application equipment after use to avoid injury to sensitive plants. To avoid injury to sensitive species, keep traffic out of treated areas until sprays have dried; irrigate soil lightly to move Torocity from turf foliage before resuming normal irrigation.

Turfgrass Use Restrictions:

- **DO NOT** overspray or allow spray to drift to ornamentals or flower beds and gardens. Roses and daylilies are particularly sensitive to Torocity.
- **DO NOT** apply more than 8 fl. oz./A (0.25 lb. AI/A) in a single application.
- **DO NOT** apply more than 16 fl. oz. (or 0.50 lb. mesotrione per acre per year) per acre per year.
- **DO NOT** make more than 4 applications of Torocity per year when using reduced application rates.
- If multiple applications are made, they must be made no closer than 14 days apart.
- **DO NOT** plant any crop other than turfgrass for 18 months post-application of Torocity to avoid turfgrass injury.
- **DO NOT** apply organophosphate or carbamate insecticides within 7 days of applying Torocity.
- **DO NOT** apply Torocity through any type of irrigation system.
- **DO NOT** make aerial applications.
- **DO NOT** use treated clippings to mulch trees or vegetable/flower gardens.
- **DO NOT** apply this product on Bentgrass, *Poa annua*, kikuyugrass, zoysiagrass, seashore paspalum, and bermudagrass plant injury is unacceptable. Maintain a 5-foot buffer between treated areas and bentgrass or *Poa annua* greens.
- **DO NOT** apply over the top of exposed roots of trees and ornamentals.
- **DO NOT** use on golf course putting greens; maintain a minimum of a 5-foot buffer between putting greens and treated areas.
- Residential Lawns: **DO NOT** make broadcast applications for pre- and post-emergent weed control unless the home lawn is being reseeded and/or renovated as whitening of some turfgrasses may occur.

Tolerant Turfgrass Species

The grasses listed below are tolerant to Torocity at the specified rates.

Species	Application Rate (fl. oz. per Acre)
Buffalograss (<i>Buchloe dactyloides</i>) Centipedegrass (<i>Eremochloa ophiuroides</i>) Kentucky bluegrass (<i>Poa pratensis</i>) Tall fescue (<i>Festuca arundinacea</i>)	5 - 8 (0.155 - 0.25 lb. AI/A)
Fine fescue* (creeping red, chewings and hard) <i>Festuca</i> spp. Perennial ryegrass* (<i>Lolium perenne</i>)	5 (0.155 lb. AI/A)
St. Augustinegrass* (grown for sod) (<i>Stenotaphrum secundatum</i>)	4 (0.125 lb. AI/A)

* See additional rate instructions below.

RESISTANCE MANAGEMENT

There is no known resistance to Torocity and no known instances of cross-resistance between the active ingredient in Torocity (mesotrione, a Group 27 HPPD inhibitor) and other classes of herbicides. Do not apply less than the specified label instructions and rotate to a herbicide product with a different mode of action to prevent the occurrence of weed resistance, and the maximum number of applications and/or rate of product has been met and weeds are not controlled.

Suspected herbicide-resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

To minimize the occurrence of resistant biotypes, observe the following weed management practices:

- Scout application site before and after herbicide applications.
- Start with a clean application site, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small.
- Add other herbicides (e.g. a selective and/or a residual herbicide) and cultural practices (e.g. tillage or crop rotation) where appropriate.
- Utilize the specified label rate for the most difficult to control weed in your field. Avoid tank mixtures with other herbicides that reduce this product's efficacy (through antagonism), or tank mixture directions that encourage application rates of this product below the label directions.
- Control weed escapes and prevent weeds from setting seeds.
- Clean equipment before moving from field to field to minimize the spread of weed seed or plant parts.
- Report any incidence of repeated non-performance of this product on a particular weed to local extension specialists, certified crop advisors, or your Atticus, LLC representative.
- Contact your local sales representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective mechanisms of actions for each target weed.

APPLICATION INSTRUCTIONS

Pre-emergence Applications:

Apply 4 - 8 fl. oz. (0.125 - 0.25 lb. AI) of **Torocity** per acre in at least 30 gallons of water per acre before seeds germinate and as close to seed germination as possible. Combine this product with another pre-emergence herbicide including Barricade 65WG (EPA Reg. No. 100-834, proflaminate) for extended control of crabgrass and foxtail.

Pre-emergence Application Precautions:

Torocity is most effective on established turf when applied post-emergence unless it is combined with another soil active herbicide.

Pre-emergence Application Restrictions:

- **DO NOT** exceed 5 fl. oz. (0.155 lb. AI) per acre per application to perennial ryegrass, fine fescues, or mixed stands that consist of >50% perennial ryegrass and/or fine fescue.
- St. Augustinegrass sod: **DO NOT** exceed 4 fl. oz. (0.125 lb. AI) per acre.

Application to New Seedlings/New Lawns

Apply 5 - 8 fl. oz. (0.155 - 0.25 lb. AI) **Torocity** per acre in at least 30 gallons of water per acre before seeding or after seeding of turfgrass species listed below, except fine fescue, as application to fine fescue can reduce grass density. **Torocity** can be effectively used on grass seed blends that contain <20% by weight hard/fine fescue. For optimal control, apply at grass seeding or as close to seeding as possible.

New Seedlings/New Lawns Restrictions:

- **DO NOT** spray on newly germinated turfgrass. Delay treatment until grass has been mowed 2 - 4 times and/or 4 weeks after emergence (whichever is longer).

Post-emergence Application Instructions:

Apply 4 - 8 fl. oz. (0.125 - 0.25 lb. AI) of **Torocity** per acre in at least 30 gallons of water per acre with a NIS surfactant. Make a repeat application 2 - 3 weeks later for optimal weed control. Apply to young, actively growing weeds.

Post-emergence Application Precautions:

Moisture stress and application to mature weeds can reduce herbicide efficacy.

Bentgrass (*Agrostis* spp.)/Nimblewill (*Muhlenbergia schreberi*) Treatment:

Apply 5 fl. oz. (0.155 lb. AI) **Torocity** per acre in at least 30 gallons of water per acre combined with a NIS surfactant at 2 - 3 week intervals for a maximum of 3 applications. For optimal weed control, apply in late summer/early fall just prior to new growth.

St. Augustinegrass (Sod Uses Only) and Centipedegrass Treatment:

Apply to established turf ONLY.

St. Augustinegrass (Sod Uses Only) and Centipedegrass Restrictions:

- **DO NOT** exceed 4 fl. oz. (0.125 lb. AI) **Torocity** per application if tank mixing with atrazine or simazine.
- **DO NOT** exceed 0.5 lb. atrazine or simazine active ingredient. See atrazine/simazine labels for precautions and restrictions.

Dormant Bermudagrass Application Only:

Apply 5 fl. oz. (0.155 lb. AI) per acre of **Torocity** to control winter weeds listed in the **WEEDS CONTROLLED** table below. Make a repeat application 2 - 3 weeks later. Application of **Torocity** to semi-dormant turf will cause bermudagrass whitening.

Spot Applications of this Product

Spray Mix	Application Rate	Rate of this Product	Rate of NIS Adjuvant
2 gallons	1 gallon per 1,000 sq. ft.	1 teaspoon	3 teaspoons

Spot Application Restrictions:

- **DO NOT** apply more than 16 fl. oz. (0.50 lb. AI) of **Torocity** per acre per year.

WEEDS CONTROLLED USING PRE-EMERGENCE APPLICATION

Apply **Torocity** with a grass pre-emergence herbicide including Barricade 65WG (EPA Reg. No. 100-834, proflaminate) herbicide, except when used to control weeds in new seedings. **Torocity** will control the following weeds using pre-emergence application:

Common Name	Scientific Name
Barnyardgrass	<i>Echinochloa crus-galli</i>
Bentgrass (Creeping)	<i>Agrostis stolonifera</i>
Bluegrass (Annual)*	<i>Poa annua*</i>
Buckhorn Plantain	<i>Plantago lanceolata</i>
Carpetweed	<i>Mollugo verticillata</i>
Chickweed (Common)	<i>Stellaria media</i>
Chickweed (Mouseear)	<i>Cerastium vulgatum</i>
Clover (Large Hop)	<i>Trifolium aureum</i>
Clover (White)	<i>Trifolium repens</i>
Crabgrass (Large)	<i>Digitaria sanguinalis</i>
Crabgrass (Smooth)	<i>Digitaria ischaemum</i>
Crabgrass (Southern)	<i>Digitaria ciliaris</i>
Foxtail (Yellow)	<i>Setaria glauca</i>
Galinsoga	<i>Galinsoga ciliata</i>
Lambsquarters	<i>Chenopodium album</i>
Pigweed (Redroot)	<i>Amaranthus retroflexus</i>
Pigweed (Smooth)	<i>Amaranthus hybridus</i>

(continued)

**WEEDS CONTROLLED USING
PRE-EMERGENCE APPLICATION (continued)**

Common Name	Scientific Name
Purslane (Common)	<i>Portulaca oleracea</i>
Shepherd's Purse	<i>Capsella bursa-pastoris</i>
Smartweed (Pale)	<i>Polygonum lapathifolium</i>
Smartweed (Pennsylvania)	<i>Polygonum pennsylvanicum</i>
Speedwell (Persian)	<i>Veronica persica</i>
Speedwell (Purslane)	<i>Veronica peregrine</i>
Wild Carrot	<i>Daucus carota</i>

*Suppression only.

**WEEDS CONTROLLED USING
POST-EMERGENCE APPLICATION**

Make a second application of **Torocity 2** - 3 weeks after initial treatment. For optimal control add a NIS-type surfactant and apply to young, actively growing weeds. **Torocity** will control the following weeds using post-emergence application:

Common Name	Scientific Name
Barnyardgrass	<i>Echinochloa crus-galli</i>
Bentgrass (Creeping)	<i>Agrostis stolonifera</i>
Buckhorn Plantain	<i>Plantago lanceolata</i>
Carpetweed	<i>Mollugo verticillata</i>
Chickweed (Common)	<i>Stellaria media</i>
Chickweed (Mouseear)	<i>Cerastium vulgatum</i>
Clover (Large Hop)	<i>Trifolium aureum</i>
Clover (White)	<i>Trifolium repens</i>
Crabgrass (Large)*	<i>Digitaria sanguinalis*</i>
Crabgrass (Smooth)*	<i>Digitaria ischaemum*</i>
Crabgrass (Southern)*	<i>Digitaria ciliaris*</i>
Curly Dock	<i>Rumex crispus</i>
Dandelion (Catsear)	<i>Hypochaeris radicata</i>
Dandelion (Common)	<i>Taraxacum officinale</i>
Florida Betony	<i>Stachys floridana</i>
Florida Pusley	<i>Richardia scabra</i>
Foxtail (Yellow)	<i>Setaria glauca</i>
Galinsoga	<i>Galinsoga ciliata</i>
Goosegrass*	<i>Eleusine indica*</i>
Ground Ivy	<i>Glechoma hederacea</i>
Healall	<i>Prunella vulgaris</i>

(continued)

**WEEDS CONTROLLED USING
POST-EMERGENCE APPLICATION (continued)**

Common Name	Scientific Name
Henbit	<i>Lamium amplexicaule</i>
Lambsquarters (Common)	<i>Chenopodium album</i>
Lawn Burweed	<i>Soliva sessilis</i>
Lovegrass (Tufted)	<i>Eragrostis pectinacea</i>
Marestail	<i>Conyza canadensis</i>
Nimblewill	<i>Muhlenbergia schreberi</i>
Nutsedge (Yellow)	<i>Cyperus esculentus</i>
Oxalis	<i>Oxalis stricta</i>
Pigweed (Redroot)	<i>Amaranthus retroflexus</i>
Pigweed (Smooth)	<i>Amaranthus hybridus</i>
Purslane (Common)	<i>Portulaca oleracea</i>
Shepherd's Purse	<i>Capsella bursa-pastoris</i>
Smartweed (Pale)	<i>Polygonum lapathifolium</i>
Smartweed (Pennsylvania)	<i>Polygonum pennsylvanicum</i>
Sowthistle	<i>Sonchus oleraceus</i>
Swinecress	<i>Coronopus didymus</i>
Thistle (Canada)	<i>Cirsium arvense</i>
Verbena	<i>Verbena hastata</i>
Wild Carrot	<i>Daucus carota</i>
Wild Violet	<i>Viola pratincola</i>
Windmillgrass	<i>Chloris verticillata</i>

*For optimal control, apply to less than 4 tiller crabgrass and goosegrass.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Keep container tightly closed when not in use. Do not store near seed, fertilizers, or foodstuffs. Can be stored at temperatures as low as minus 20°F. Keep away from heat and flame.

Pesticide Disposal: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Open dumping is prohibited.

Container Handling ≤ 5 Gallons: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

STORAGE AND DISPOSAL *(continued)*

Container Handling > 5 Gallons: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Refillable Containers: Refillable container. Refill this container with pesticide only. Do not use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of Atticus, LLC. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Atticus, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, neither Atticus, LLC, the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

Torocity™ is a trademark of Atticus, LLC
20210223ap1